

Table S5. Pairwise niche overlap statistics between four *Drusus* species in the Western Balkans region. Three niche overlap statistics (Schoener's D (D), the I-statistic (I), and the relative rank statistic (RR)) were calculated based on Maxent models for present-day climatic conditions and two LGM scenarios (MIROC and CCSM).

CURRENT	<i>D. discolor</i>			<i>D. krusniki</i>			<i>D. schmidi</i>		
	D	I	RR	D	I	RR	D	I	RR
<i>D. croaticus</i>	0.131	0.335	0.713	0.149	0.396	0.590	0.283	0.587	0.661
<i>D. discolor</i>	-	-	-	0.190	0.428	0.577	0.365	0.669	0.679
<i>D. krusniki</i>				-	-	-	0.494	0.790	0.854
MIROC	<i>D. discolor</i>			<i>D. krusniki</i>			<i>D. schmidi</i>		
	D	I	RR	D	I	RR	D	I	RR
<i>D. croaticus</i>	0.105	0.289	0.732	0.075	0.191	0.709	0.220	0.437	0.756
<i>D. discolor</i>	-	-	-	0.313	0.578	0.704	0.485	0.766	0.715
<i>D. krusniki</i>				-	-	-	0.453	0.753	0.862
CCSM	<i>D. discolor</i>			<i>D. krusniki</i>			<i>D. schmidi</i>		
	D	I	RR	D	I	RR	D	I	RR
<i>D. croaticus</i>	0.036	0.128	0.762	0.017	0.066	0.663	0.023	0.092	0.756
<i>D. discolor</i>	-	-	-	0.490	0.765	0.778	0.684	0.913	0.820
<i>D. krusniki</i>				-	-	-	0.577	0.843	0.854